

| | | Contraceptive prevalence rate | Skilled attendant at delivery | C-section rate | Antenatal care (4+ visits) | Early initiation of breastfeeding | Postnatal care for babies born at home | Postnatal care for all babies* | Family planning needs satisfied | Improved water sources | DPT3 vaccine | Measles vaccine | BCG vaccine | Vitamin A administration | ORT and continued feeding | Care seeking for pneumonia | Insecticide-treated use by children | Co-coverage (6 or more) | Coverage gap | Mean coverage |
|---------------------|------------|-------------------------------|-------------------------------|----------------|----------------------------|-----------------------------------|----------------------------------------|--------------------------------|---------------------------------|------------------------|--------------|-----------------|-------------|--------------------------|---------------------------|----------------------------|-------------------------------------|-------------------------|--------------|---------------|
| Child's sex | Male | NA | 61,7% | 14,6% | 58,4% | 60,3% | 4,2% | 60,0% | NA | NA | 70,7% | 65,0% | 94,0% | 60,2% | 56,6% | 50,2% | ND | 12,0% | NA | NA |
| | Female | NA | 60,3% | 14,6% | 57,8% | 61,5% | 5,1% | 58,3% | NA | NA | 74,1% | 64,0% | 93,3% | 62,8% | 49,9% | 46,0% | ND | 12,3% | NA | NA |
| Region of residence | Urban | 64,0% | 77,9% | 21,3% | 68,2% | 60,8% | 3,1% | 76,5% | 77,7% | 96,1% | 75,0% | 66,9% | 95,1% | 62,8% | 55,5% | 51,0% | ND | 15,4% | NA | NA |
| | Rural | 48,0% | 38,7% | 5,8% | 42,5% | 61,1% | 5,4% | 36,3% | 61,2% | 76,2% | 68,6% | 60,9% | 91,7% | 59,6% | 51,7% | 44,4% | ND | 7,8% | NA | NA |
| Wealth quintiles | Q1 | 41,8% | 26,7% | 2,7% | 34,5% | 59,5% | 4,2% | 24,8% | 52,5% | 62,1% | 65,5% | 63,1% | 91,1% | 58,0% | 47,9% | 39,7% | ND | 5,8% | 50,4% | 49,6% |
| | Q2 | 53,9% | 49,9% | 6,8% | 48,8% | 62,7% | 6,5% | 48,8% | 68,2% | 89,8% | 70,9% | 58,0% | 90,8% | 60,8% | 53,9% | 53,9% | ND | 9,7% | 39,0% | 61,0% |
| | Q3 | 58,4% | 70,1% | 13,6% | 59,3% | 60,5% | 3,5% | 66,9% | 71,1% | 94,7% | 66,3% | 62,0% | 92,8% | 61,6% | 54,5% | 49,3% | ND | 13,1% | 35,1% | 64,9% |
| | Q4 | 64,6% | 87,9% | 21,5% | 71,3% | 63,2% | 2,8% | 86,1% | 78,1% | 97,2% | 81,6% | 70,4% | 97,9% | 63,2% | 56,8% | 47,1% | ND | 17,2% | 26,9% | 73,1% |
| | Q5 | 70,9% | 98,6% | 47,1% | 88,8% | 57,9% | 7,0% | 98,2% | 87,2% | 98,9% | 85,5% | 74,4% | 99,1% | 66,6% | 64,0% | 56,9% | ND | 20,6% | 18,1% | 81,9% |
| Maternal education | A | 33,6% | 30,1% | 4,9% | 32,2% | 57,3% | 3,0% | 28,1% | 51,7% | NO | 58,5% | 60,9% | 89,2% | 58,5% | 52,2% | 40,0% | ND | 4,9% | NA | NA |
| | B | 54,0% | 48,5% | 7,6% | 45,9% | 60,7% | 4,6% | 46,3% | 66,6% | NO | 65,4% | 59,1% | 91,5% | 58,5% | 51,4% | 44,7% | ND | 8,8% | NA | NA |
| | C | 64,8% | 78,9% | 17,6% | 68,8% | 65,0% | 8,3% | 78,2% | 76,4% | NO | 81,0% | 69,7% | 95,7% | 65,0% | 53,7% | 56,6% | ND | 17,3% | NA | NA |
| | D | 71,2% | 94,1% | 35,8% | 85,4% | 59,7% | 2,9% | 92,3% | 85,1% | NO | 87,7% | 74,8% | 99,0% | 67,5% | 62,7% | 53,4% | ND | 20,5% | NA | NA |
| Country region | chiquisaca | 50,6% | 53,9% | 8,6% | 62,0% | 59,1% | 8,6% | 54,8% | 62,4% | 71,9% | 88,0% | 76,6% | 98,9% | 74,9% | 59,2% | 47,9% | ND | 11,2% | NA | NA |
| | la paz | 56,1% | 44,9% | 8,1% | 46,1% | 60,0% | 2,7% | 42,4% | 71,7% | 92,6% | 60,5% | 51,6% | 88,9% | 57,1% | 50,5% | 35,8% | ND | 9,5% | NA | NA |
| | cochabamba | 56,7% | 62,1% | 17,6% | 55,9% | 63,7% | 3,5% | 60,1% | 70,3% | 80,8% | 71,2% | 63,4% | 96,0% | 59,6% | 56,2% | 42,5% | ND | 11,1% | NA | NA |
| | oruro | 48,7% | 58,4% | 9,3% | 56,7% | 52,7% | 7,7% | 57,5% | 63,0% | 97,9% | 74,0% | 65,8% | 94,8% | 61,1% | 53,3% | 49,8% | ND | 10,7% | NA | NA |
| | potosi | 48,6% | 42,8% | 6,8% | 50,9% | 49,7% | 6,7% | 40,7% | 61,7% | 80,5% | 73,5% | 65,3% | 94,3% | 60,5% | 55,5% | 52,8% | ND | 10,7% | NA | NA |
| | tarija | 67,6% | 79,3% | 18,4% | 75,8% | 64,0% | 6,7% | 79,7% | 81,1% | 87,2% | 90,5% | 79,2% | 99,5% | 70,7% | 60,4% | 60,9% | ND | 16,8% | NA | NA |
| | santa cruz | 66,5% | 84,7% | 26,0% | 70,1% | 64,5% | 5,1% | 82,4% | 78,3% | 96,9% | 78,6% | 71,6% | 94,3% | 60,6% | 50,6% | 54,1% | ND | 16,5% | NA | NA |
| | beni | 62,1% | 68,9% | 12,9% | 65,3% | 69,3% | 5,2% | 67,3% | 75,4% | 86,1% | 65,6% | 66,7% | 91,7% | 68,3% | 56,2% | 56,1% | ND | 12,5% | NA | NA |
| | pando | 60,5% | 69,6% | 12,4% | 75,7% | 73,3% | 3,1% | 67,6% | 73,4% | 65,2% | 76,2% | 77,8% | 90,1% | 74,2% | 64,6% | 61,0% | ND | 9,6% | NA | NA |
| Overall | | 58,4% | 61,0% | 14,6% | 58,1% | 60,9% | 4,6% | 59,2% | 72,0% | 88,6% | 72,4% | 64,5% | 93,7% | 61,4% | 53,7% | 48,3% | ND | 12,2% | 35,4% | 64,6% |
| Q5/Q1 ratio | | 1,70 | 3,69 | 17,36 | 2,58 | 0,97 | 1,68 | 3,95 | 1,66 | 1,59 | 1,31 | 1,18 | 1,09 | 1,15 | 1,34 | 1,43 | ND | 3,54 | 0,36 | 1,65 |
| Q5-Q1 difference | | 29,2 | 71,9 | 44,4 | 54,3 | -1,7 | 2,8 | 73,4 | 34,7 | 36,7 | 20,0 | 11,2 | 7,9 | 8,6 | 16,1 | 17,2 | ND | 14,8 | -32,3 | 32,3 |
| Slope index | | 32,7 | 89,7 | 44,3 | 62,1 | -0,1 | -0,3 | 90,2 | 37,3 | 39,7 | 22,4 | 14,9 | 10,6 | 8,9 | 15,6 | 12,4 | ND | 17,6 | NA | NA |
| Concentration index | | 9,4 | 24,4 | 48,4 | 17,9 | 0,2 | 0,0 | 25,2 | 8,8 | 7,8 | 4,9 | 3,8 | 1,8 | 2,3 | 4,1 | 4,9 | ND | 23,8 | NA | NA |




Fig 1. Mean coverage of selected interventions and corresponding coverage gap (how much is needed to reach universal coverage), by wealth quintile.

| | Antenatal care (4+ visits) | Skilled attendant at delivery | Care seeking for pneumonia | ORT and continued feeding | BCG vaccine | DPT3 vaccine | Measles vaccine | Family planning needs satisfied |
|------------------|----------------------------|-------------------------------|----------------------------|---------------------------|-------------|--------------|-----------------|---------------------------------|
| Poorest quintile | 34,5 | 26,7 | 39,7 | 47,9 | 91,1 | 65,5 | 63,1 | 52,5 |
| Richest quintile | 54,3 | 71,9 | 17,2 | 16,1 | 7,9 | 20,0 | 11,2 | 34,7 |
| Richest quintile | 88,8 | 98,6 | 56,9 | 64,0 | 99,1 | 85,5 | 74,4 | 87,2 |




Fig 2. Coverage levels in the poorest and richest quintiles for selected interventions along the continuum of care.

| Co-coverage interv | Wealth quintile | | | | |
|--------------------|-----------------|-------|-------|-------|-------|
| | Q1 | Q2 | Q3 | Q4 | Q5 |
| 0 | 0,112 | 0,023 | 0,006 | 0,002 | 0,000 |
| 1 | 0,283 | 0,191 | 0,124 | 0,042 | 0,006 |
| 2 | 0,262 | 0,276 | 0,227 | 0,191 | 0,076 |
| 3 | 0,143 | 0,207 | 0,258 | 0,261 | 0,282 |
| 4 | 0,085 | 0,125 | 0,150 | 0,202 | 0,270 |
| 5 | 0,057 | 0,081 | 0,105 | 0,131 | 0,160 |
| 6 | 0,036 | 0,036 | 0,056 | 0,063 | 0,062 |
| 7 | 0,015 | 0,042 | 0,047 | 0,067 | 0,097 |
| 8 | 0,006 | 0,019 | 0,028 | 0,042 | 0,047 |
| 9 | | | | | |
| Total | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 |




Fig 3. Percentage of children aged 1-4 years according to the number of key child-survival interventions received, by wealth quintile.