Countdown to 2015
Maternal, Newborn & Child Survival

Bolivia

March 2012

DEMOGRAPHICS

Total population (000)	9,930	(2010)
Total under-five population (000)	1,235	(2010)
Births (000)	263	(2010)
Birth registration (%)	74	(2001)
Total under-five deaths (000)	14	(2010)
Neonatal deaths: % of all under-5 deaths	44	(2010)
Neonatal mortality rate (per 1000 live births)	23	(2010)
Infant mortality rate (per 1000 live births)	42	(2010)
Stillbirth rate (per 1000 total births)	17	(2009)
Total maternal deaths	470	(2008)
Lifetime risk of maternal death (1 in N)	150	(2008)
Total fertility rate (per woman)	3.3	(2010)
Adolescent birth rate (per 1000 women)	89	(2006)

Under-five mortality rate

Deaths per 1000 live births



Source: IGME 2011

Maternal mortality ratio

Deaths per 100,000 live births



Source: MMEIG 2010

Note: MDG target calculated by Countdown to 2015

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (%, of total ARVs) 1 (2010)



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization

- Percent of children immunized against measles
- Percent of children immunized with 3 doses DTP
- Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

- Percent children <5 years with suspected pneumonia taken to appropriate health provider
- Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	1	(2008)
Low birthweight incidence (moderate and severe, %)	6	(2008)
Early initiation of breastfeeding (within 1 hr of birth, %)	64	(2008)
Introduction of solid, semi-solid/soft foods (%)	83	(2008)
Vitamin A two dose coverage (%)	24	(2010)

Underweight and stunting prevalence

- Percent children <5 years who are underweight
- Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed