





DEMOGRAPHICS


Total population (000)	1,728	(2010)
Total under-five population (000)	287	(2010)
Births (000)	66	(2010)
Birth registration (%)	55	(2005-2006)
Total under-five deaths (000)	6	(2010)
Neonatal deaths: % of all under-5 deaths	34	(2010)
Neonatal mortality rate (per 1000 live births)	31	(2010)
Infant mortality rate (per 1000 live births)	57	(2010)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	250	(2008)
Lifetime risk of maternal death (1 in N)	49	(2008)
Total fertility rate (per woman)	4.9	(2010)
Adolescent birth rate (per 1000 women)	104	(2000)



Note: MDG target calculated by Countdown to 2015


MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care



Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
▭ Uncertainty range around the estimate



EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH


Immunization

■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2005-2006)
Low birthweight incidence (moderate and severe, %) 20 (2005-2006)
Early initiation of breastfeeding (within 1 hr of birth, %) 48 (2005-2006)
Introduction of solid, semi-solid/soft foods (%) -
Vitamin A two dose coverage (%) 100 (2010)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed

