




DEMOGRAPHICS

Total population (000)	46,218	(2011)
Total under-five population (000)	8,273	(2011)
Births (000)	1,913	(2011)
Birth registration (%)	16	(2010)
Total under-five deaths (000)	122	(2011)
Neonatal deaths: % of all under-5 deaths	40	(2011)
Neonatal mortality rate (per 1000 live births)	25	(2011)
Infant mortality rate (per 1000 live births)	45	(2011)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	8,500	(2010)
Lifetime risk of maternal death (1 in N)	38	(2010)
Total fertility rate (per woman)	5.5	(2011)
Adolescent birth rate (per 1000 women)	128	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

Percent HIV+ pregnant women receiving ARVs for PMTCT


Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



DHS 2010


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	5	(2010)	Early initiation of breastfeeding (within 1 hr of birth, %)	49	(2010)
Low birthweight incidence (moderate and severe, %)	8	(2010)	Introduction of solid, semi-solid/soft foods (%)	92	(2010)
			Vitamin A two dose coverage (%)	97	(2011)

Underweight and stunting prevalence

Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed

