




DEMOGRAPHICS

Total population (000)	8,445 (2006)
Total under-five population (000)	1,507 (2006)
Births (000)	371 (2006)
Birth registration (%)	3 (2006)
Under-five mortality rate (per 1000 live births)	145 (2006)
Infant mortality rate (per 1000 live births)	90 (2006)
Neonatal mortality rate (per 1000 live births)	49 (2006)
Total under-five deaths (000)	54 (2006)
Maternal mortality ratio (per 100,000 live births)	1,400 (2005)
Lifetime risk of maternal death (1 in N)	12 (2005)
Total maternal deaths	5,200 (2005)

Under-five mortality rate


Deaths per 1000 live births



Source: UNICEF, 2006


Causes of under-five deaths

Globally more than one third of child deaths are attributable to undernutrition



Source: WHO, 2006

Causes of neonatal deaths




Source: Lawn JE, Cousens SN for CHERG (Nov 2006)

MATERNAL AND NEWBORN HEALTH

Unmet need for family planning (%)	---
Antenatal visits for woman (4 or more visits, %)	---
Intermittent preventive treatment for malaria (%)	1 (2006)
C-section rate (total, urban, rural, %)	---
Early initiation of breastfeeding (within 1 hr of birth, %)	26 (2006)
Postnatal visit for baby (within 2 days for home births, %)	---


Causes of maternal deaths

Regional estimates for Africa, 1997-2002



Source: Khan, Khalid S., et al., Lancet 2006;367:1066-74

Coverage along the continuum of care



Source: DHS, MICS, Other NS

*See Annex for indicator definition


INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

NUTRITION

Stunting prevalence (moderate and severe, %)	42 (2006)	Complementary feeding rate (6-9 months, %)	15 (2006)
Wasting prevalence (moderate and severe, %)	13 (2006)	Low birthweight incidence (%)	11 (2006)

Underweight prevalence


Percent children < 5 years underweight for age*



*Based on 2006 WHO reference population


Exclusive breastfeeding

Percent infants < 6 months exclusively breastfed



Vitamin A supplementation


Percent children 6-59 months receiving vitamin A doses



Source: UNICEF


Antenatal care

Percent women aged 15-49 years attended at least once by a skilled health provider during pregnancy




Skilled attendant at delivery

Percent live births attended by skilled health personnel



Neonatal tetanus protection

Percent of newborns protected against tetanus




Source: WHO/UNICEF

CHILD HEALTH

Immunization


Percent of children immunised against measles
Percent of children immunised with 3 doses DPT
Percent of children immunised with 3 doses Hib



Source: WHO/UNICEF

Malaria prevention


Percent children < 5 years sleeping under ITNs



Source: MICS

Prevention of mother to child transmission of HIV


Percent HIV+ pregnant women receiving ARVs for PMTCT



Source: Other NS

Diarrhoeal disease treatment


Percent children < 5 years with diarrhoea receiving oral rehydration therapy or increased fluids, with continued feeding



Source: MICS

Malaria treatment


Percent febrile children < 5 years using antimalarials



Source: MICS

Pneumonia treatment

Percent children < 5 years with suspected pneumonia taken to appropriate health provider
Percent children < 5 years with suspected pneumonia receiving antibiotics




Source: MICS

WATER AND SANITATION


Water

Percent population using improved drinking water sources



Sanitation

Percent population using improved sanitation facilities



Source: WHO/UNICEF JMP, 2006

POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	---
Community treatment of pneumonia with antibiotics	---
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	No
Midwives be authorised to administer a core set of life saving interventions	---
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	---

SYSTEMS

Per capita total expenditure on health (US\$)	---
General government expenditure on health as % of total government expenditure (%)	---
Out-of-pocket expenditure as % of total expenditure on health (%)	---
Density of health workers (per 1000 population)	0.2 (1997)
Official Development Assistance to child health per child (US\$)	4 (2005)
Official Development Assistance to maternal and neonatal health per live birth (US\$)	4 (2005)
National availability of Emergency Obstetric Care services (% of recommended minimum)	56 (2005)

EQUITY

Coverage gap by wealth quintile

Coverage gap (%)	No data
Ratio poorest/wealthiest	No data
Difference poorest-wealthiest (%)	No data