




DEMOGRAPHICS

Total population (000)	10,124	(2011)
Total under-five population (000)	1,246	(2011)
Births (000)	266	(2011)
Birth registration (%)	81	(2005-2006)
Total under-five deaths (000)	19	(2011)
Neonatal deaths: % of all under-5 deaths	36	(2011)
Neonatal mortality rate (per 1000 live births)	25	(2011)
Infant mortality rate (per 1000 live births)	53	(2011)
Stillbirth rate (per 1000 total births)	15	(2009)
Total maternal deaths	940	(2010)
Lifetime risk of maternal death (1 in N)	83	(2010)
Total fertility rate (per woman)	3.3	(2011)
Adolescent birth rate (per 1000 women)	69	(2003)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH


Coverage along the continuum of care



Source: DHS, MICS, Other NS
* See Annex/website for indicator definition


Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)



Source: UNICEF/UNAIDS/WHO

EQUITY


Socioeconomic inequities in coverage



DHS 2005-2006
Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH


Immunization



Source: WHO/UNICEF

Pneumonia treatment


Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)	10 (2005-2006)	Early initiation of breastfeeding (within 1 hr of birth, %)	44 (2005-2006)
Low birthweight incidence (moderate and severe, %)	25 (2005-2006)	Introduction of solid, semi-solid/soft foods (%)	90 (2005-2006)
		Vitamin A two dose coverage (%)	36 (2011)

Underweight and stunting prevalence



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

